

Associations

Phasey bean can grow in association with summer grasses to provide early summer grazing. Companion species include lespedeza species (*Kummerowia stipulacea*, *Kummerowia striata*, *Lepedeza cuneata*), *Chloris gayana*, *Dichanthium aristatum*, several *Paspalum* species (*Paspalum dilatatum* and *Paspalum plicatulum*) and Para grass (*Brachiaria mutica*). Association with Guinea grass (*Megathyrsus maximus*), broadleaf setaria (*Setaria sphacelata* var. *splendida*), and scrobbic is successful in well-drained conditions. Warm season legumes that are valuable companions include American vetch (*Aeschynomene americana*), white clover (*Trifolium repens*) and *Desmodium heterocarpon* (Cook et al., 2005). Phasey bean may also be sown in wheat or maize fields (Asongwed-Awa et al., 2002; Muldoon, 1984).

In poorly drained soils or irrigated areas the seeds should be sown into the top of planting ridges (FAO, 2012). For instance, in association with Para grass, phasey bean is sown on the tops of ridges and 2 to 3 months later the grass is sown in the ridges where irrigation water enters. Phasey bean is also frequently grown in Zimbabwe and Sudan, and also in Mali in irrigated cotton rotations where it is used for grazing and for hay (Göhl, 1982).

Pasture

Phasey bean should not be grazed heavily or continuously as this may hamper its viability. Light grazing and/or rotational grazing (leaving at least 10-15 cm growth) followed by a 6 to 8 week rest period in the growing season is adequate to help phasey bean regeneration (FAO, 2012; Cook et al., 2005). As a short-lived legume, *Macroptilium lathyroides* can help to establish grasses such as *Desmodium heterocarpon* which is persistent under grazing, but not easily established (Aiken et al., 1991b). An overseeding of *Paspalum notatum* with a mixture of *Desmodium heterocarpon*, *Aeschynomene americana*, and *Macroptilium lathyroides* was tested with yearling steers. During the establishment period, phasey bean and *Aeschynomene* provided high quality forage, but contributed negligibly during the second year (Aiken et al., 1991a).

Hay and silage

Phasey bean makes good quality hay provided that it is cut and handled early enough to preserve a maximum of leaf material (FAO, 2012). It also makes excellent silage in combination with Columbus grass (*Sorghum x alnum*) or alone: it proved to have higher fermentation potential than Guinea grass (Imura et al., 2001; Robertson, 1971 cited by FAO, 2012).

Environmental impact

N-fixing legume, cover crop and biodiversity

Phasey bean is an N-fixing legume that nodulates freely with native rhizobia, making seed inoculation unnecessary. Its association with scrobbic (*Paspalum commersonii*) increased forage grass yield by 77% and was found equivalent to the application of 800 kg/ha sulphate of ammonia. After a 3-year cultivation, phasey bean increased soil N content by 10-15% at a depth of 60-90 cm (FAO, 2012). Phasey bean is often cited as a potential cover crop, especially under flooded conditions (Werner et al., 2005). It can also be used in wildlife-food plantings to provide seeds for quail and forage for deer (Newman et al., 2002).

Nematode sensitivity

Phasey bean hosts knot-root nematode and is very sensitive to them. This can become a problem in nematode control (Rich et al., 2010).

Weed

Phasey bean competes with weeds and is seldom referred to as a weed. However, its ability to twin may become a problem in citrus plantations (Barnes et al., 2007).

Datasheet citation

Heuzé V., Tran G., Giger-Reverdin S., Bastianelli D., Lebas F., 2015. *Phasey bean (Macroptilium lathyroides)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/627> Last updated on September 30, 2015, 17:30


English correction by Tim Smith (Animal Science consultant) and Hélène Thiollot (AFZ)

Image credits

- Forest and Kim Starr ● Forest and Kim Starr



Automatic translation

 Sélectionner une langue ▼

Feed categories

All feeds

Forage plants

- Cereal and grass forages
- Legume forages
- Forage trees
- Aquatic plants
- Other forage plants

Plant products/by-products

- Cereal grains and by-products
- Legume seeds and by-products
- Oil plants and by-products
- Fruits and by-products
- Roots, tubers and by-products
- Sugar processing by-products
- Plant oils and fats
- Other plant by-products

Feeds of animal origin

- Animal by-products
- Dairy products/by-products
- Animal fats and oils
- Insects

Other feeds

- Minerals
- Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- Literature databases
- Feeds and plants databases
- Organisations & networks
- Books
- Journals

Phasey bean (Macroptilium lathyroides)

- Description
- Nutritional aspects
- Nutritional tables
- References

Nutritional attributes

Information about the composition of phasey bean forage and seeds is scarce. Phasey bean forage is rich in protein (up to 27% DM) though lower values (under 15% DM) have been recorded. Fibre content is relatively high (average ADF content of 25% DM) but is quite variable. Composition depends on maturity: between 26 and 106 days, the protein content of the leaves decreased from 30 to 20% while NDF increased from 50 to 60%. The ADF content of the leaves rose from 35 to 45% between 26 and 50 days and decreased back to its original value at 106 days (Nakanishi et al., 1993). Other factors influence the composition: flooding increases the amount of structural carbohydrates at the expense of protein content, while drought has the inverse effect (Nagashiro et al., 1995). Longer cutting intervals also had a depressive effect on leaf protein content (Adjei et al., 1985).

Potential constraints

Forage

No evidence of toxicity has been found in cattle or in horses (Cook et al., 2005).

Seeds

Phasey bean seeds contain low levels of non-toxic lectin and moderate amounts of trypsin inhibitors (Grant et al., 1995). They do not require heat-treatment prior to use (Grant et al., 1991).

Ruminants

Whole plant

Fresh forage

There is limited recent information on the nutritional value of phasey bean forage. The protein content is high but is reported to be highly rumen-degradable (75%): a diet based on phasey bean forage should be supplemented with sources of rumen undegradable protein (DiCostanzo et al., 2006). Digestibility is not very high: 50-65% for *in vivo* OM digestibility measured in sheep (Milford, 1967) and about 65% for *in vitro* DM digestibility (Nasrullah et al., 2003; Nakanishi et al., 1993).

Silage

Phasey bean forage ensiled at 60 days regrowth exhibited a good fermentation pattern with a pH under 4.8 and high lactic acid fermentation. Further heat treatment (70°C) of silage did not change the amount of rumen degradable protein (about 45% of the total protein), but decreased the soluble protein content (from 19 to 8% of the protein) (Kawamoto et al., 1997).

Ensiling rice forage and phasey bean forage together improved the fermentation quality and the palatability of rice forage silages, which are lower for rice than for phasey bean (Tobisa et al., 2006; Tobisa et al., 2005). Ensiling phasey bean forage with Napier grass (*Pennisetum purpureum*) improved the nutritive value compared to Napier grass alone (Yunus et al., 2001).

Cattle

Heat treated silage of phasey bean increased the sum of rumen undegradable and indigestible protein (from 36 to 47%) (Kawamoto et al., 1997). Acid detergent insoluble N increased with heat treatment and silage DM (Tamaki et al., 2003) and was partially degradable in the post-rumen digestive tract (Tamaki et al., 2002). Dry matter and N degradability decreased as silage DM increased (Tamaki et al., 2003).

Goats

Goats grazing phasey bean pastures are able to select the leaves, which contain more protein (9% or over) and less fibre than the rest of the plant, and have a higher *in vitro* DM digestibility (75 vs. 60%). This selective grazing is made possible by the greater accessibility of the upper layers of pasture, which are younger than the less accessible lower parts (Nakanishi et al., 1993).

Seeds

Phasey bean seeds have a high protein content (27% DM) with an *in vitro* gas production similar to that of *Leucaena leucocephala*. *In sacco* DM degradation is rapid and reaches 92% after 96 hours of incubation. The seeds do not have any significant defaunating activity (Odeyinka et al., 2004).

Pigs


No information found (2012).

Poultry

No information found (2012).

Rabbits

Very few studies have been published on the utilization of phasey bean forage by rabbits. The palatability of the fresh plant



seems good: in a field study conducted in Texas, wild rabbits preferred to graze phasey bean rather than grass in a mixed culture of this legume forage and crabgrass (*Digitaria ciliaris*) ([Nguluve et al., 2004](#)).

In the only available study, phasey bean forage (whole plant, 16.8% crude protein and 40% ADF) was harvested and dried in Puerto Rico, sent to Oregon and then incorporated at 40% in a complete diet offered for 21 days to weanling New Zealand White rabbits. The growth rate obtained with this diet was identical (40.6 g/d) to that obtained with the same proportion of [alfalfa](#) hay (13.8% CP and 32% ADF). Nevertheless, crude protein digestibility was significantly reduced in the phasey bean diet when compared to that of the alfalfa diet (71 vs. 80%) ([Harris et al., 1981](#); [Cheeke et al., 1983](#)). From these results, phasey bean could be considered as a suitable forage for rabbit nutrition, but more experiments are required, particularly to determine protein availability.

Datasheet citation

Heuzé V., Tran G., Giger-Reverdin S., Bastianelli D., Lebas F., 2015. *Phasey bean (Macroptilium lathyroides)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/627> Last updated on September 30, 2015, 17:30

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

- Forest and Kim Starr
- Forest and Kim Starr



Automatic translation

Sélectionner une langue ▼

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

Description Nutritional aspects **Nutritional tables** References

Tables of chemical composition and nutritional value

- Phasey bean (*Macroptilium lathyroides*), aerial part, fresh

Avg: average or predicted value; SD: standard deviation; Min: minimum value; Max: maximum value; Nb: number of values (samples) used

Phasey bean (*Macroptilium lathyroides*), aerial part, fresh



Main analysis	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	17.8	3.5	11.8	26.2	126
Crude protein	% DM	16.7	3.6	9.6	24.8	140
Crude fibre	% DM	27.0	4.1	20.8	36.2	130
NDF	% DM	43.5	9.9	34.8	58.1	5 *
ADF	% DM	32.4	11.0	15.5	43.5	5 *
Lignin	% DM	6.2	2.8	3.7	10.5	5
Ether extract	% DM	2.8	0.7	1.8	4.5	130
Ash	% DM	8.3	1.7	5.5	12.7	132
Gross energy	MJ/kg DM	18.7				*

Minerals	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	13.1	2.9	7.8	20.3	127
Phosphorus	g/kg DM	2.2	0.5	1.2	3.8	129
Potassium	g/kg DM	16.4	4.3	10.0	27.2	125
Sodium	g/kg DM	0.4	0.3	0.1	0.6	3
Magnesium	g/kg DM	4.1	0.7	2.7	5.6	126
Manganese	mg/kg DM	99				1
Zinc	mg/kg DM	27				1
Copper	mg/kg DM	8				1

Ruminant nutritive values	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	59.8	5.8	50.3	64.7	5
Energy digestibility, ruminants	%	57.2				*
DE ruminants	MJ/kg DM	10.7				*
ME ruminants	MJ/kg DM	8.5				*
Nitrogen digestibility, ruminants	%	70.8	10.8	52.8	79.0	5

Pig nutritive values	Unit	Avg	SD	Min	Max	Nb
Energy digestibility, growing pig	%	47.7				*
DE growing pig	MJ/kg DM	8.9				*

The asterisk * indicates that the average value was obtained by an equation.

References

Barnes, 1999; Brink et al., 1988; CIRAD, 1991; Holm, 1971; Milford, 1967; Muir, 2002; Nasrullah et al., 2003

Last updated on 19/03/2013 11:55:56

Datasheet citation

Heuzé V., Tran G., Giger-Reverdin S., Bastianelli D., Lebas F., 2015. *Phasey bean (Macroptilium lathyroides)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/627> Last updated on September 30, 2015, 17:30

English correction by Tim Smith (Animal Science consultant) and H  l  ne Thiollet (AFZ)

Image credits

- Forest and Kim Starr • Forest and Kim Starr

 Share / Save   



References

Nagashiro, C. W. ; Shibata, F., 1995. Influence of flooding and drought conditions on herbage yield and quality of phasey

 Sélectionner une langue ▼

Feed categories

All feeds

Forage plants

- ▶ Cereal and grass forages
- ▶ Legume forages
- ▶ Forage trees
- ▶ Aquatic plants
- ▶ Other forage plants

Plant products/by-products

- ▶ Cereal grains and by-products
- ▶ Legume seeds and by-products
- ▶ Oil plants and by-products
- ▶ Fruits and by-products
- ▶ Roots, tubers and by-products
- ▶ Sugar processing by-products
- ▶ Plant oils and fats
- ▶ Other plant by-products

Feeds of animal origin

- ▶ Animal by-products
- ▶ Dairy products/by-products
- ▶ Animal fats and oils
- ▶ Insects

Other feeds

- ▶ Minerals
- ▶ Other products

Latin names

Plant and animal families

Plant and animal species

Resources

Broadening horizons

Literature search

Image search

Glossary

External resources

- ▶ Literature databases
- ▶ Feeds and plants databases
- ▶ Organisations & networks
- ▶ Books
- ▶ Journals

bean (*Macroptilium lathyroides* (L.) Urb.). Grassl. Sci., 41: 218-225

Nakanishi, Y. ; Tsuru, K. ; Bungo, T. ; Shimojo, M. ; Masuda, Y. ; Goto, I., 1993. Effects of growth stage and sward structure of *Macroptilium lathyroides* and *M. atropurpureum* on selective grazing and bite size in goats. Trop. Grassl., 27 (2): 108-113

Nasrullah; Niimi, M. ; Akashi, R. ; Kawamura, O., 2003. Nutritive evaluation of forage plants grown in South Sulawesi, Indonesia. Asian-Aust. J. Anim.Sci., 16 (5):693-701

Nasrullah; Niimi, M. ; Akashi, R. ; Kawamura, O., 2004. Nutritive evaluation of forage plants grown in South Sulawesi, Indonesia. II. Mineral composition. Asian-Aust. J. Anim. Sci., 17 (1): 63-67

Newman, Y. C. ; Vendramini, J. ; Blount, A. R., 2002. Minor use summer annual forage legumes. University of Florida, IFAS, SS-AGR-79

Ngulube, D. W. ; Muir, J. P. ; Wittiec, R. ; Rosiere, R. ; Butler, T. J., 2004. Yield and nutritive value of summer legumes as influenced by dairy manure compost and competition with crabgrass. Agron. J., 96 (3): 812-817

Odeyinka, S. M. ; Hector, B. L. ; Ørskov, E. R. ; Newbold, C. J., 2004. Assessment of the nutritive value of the seeds of some tropical legumes as feeds for ruminants. Livest. Res. Rural Dev., 16 (9)

Pitman, W. D. ; Kretschmer, A. E. Jr., 1984. Persistence of selected tropical pasture legumes in peninsular Florida. Agron. J., 76 (6): 993-996

Pitman, W. D. ; Portier, K. M. ; Chambliss, C. G. ; Kretschmer, A. E. Jr., 1992. Performance of yearling steers grazing bahia grass pastures with summer annual legumes or nitrogen fertiliser in subtropical Florida. Trop. Grassl., 26 (3): 206-211

Pitman, W. D., 1983. Initial comparisons of tropical legume Bahiagrass pastures and nitrogen fertilized Bahiagrass pastures in peninsular Florida. Proceedings, Soil and Crop Science Society of Florida. 1983, 42, 72-75

Rich, J. ; Brito, J. ; Ferrell, J. ; Kaur, R., 2010. Weed hosts of root-knot nematodes common to Florida. University of Florida, IFAS, ENY-060

Roberge, G. ; Toutain, B., 1999. Choix des plantes fourragères. In: Cultures fourragères tropicales, Roberge, G. ; Toutain, B. (éditeurs), Collection Repères, CIRAD, Quae Editions

Tamaki, M. ; Kawamoto, Y. ; Imura, Y. ; Motomura, T. ; Nakada, T. ; Sonoda, T., 2002. Effects of heat treatment of tropical legume phasey bean (*Macroptilium lathyroides*) silages on degradability and utilization of nitrogen in the rumen and post-ruminal digestive tracts. Grassland Science, 48 (3): 236-241

Tamaki, M. ; Kawamoto, Y. ; Motomura, T. ; Namihiro, T. ; Miyagi, T. ; Nakada, T. ; Sonoda, T., 2003. Effects of moisture content and heat treatment of tropical legume phasey bean (*Macroptilium lathyroides*) silages on degradability and utilization of nitrogen in the rumen and post-ruminal digestive tracts. Grassland Science, 49 (4): 361-366

Tobisa, M. ; Shimojo, M. ; Nakano, Y. ; Okano, K. ; Masuda, Y., 2000. Dry matter digestibility and nitrogen content of tropical forage legume *Aeschynomene americana* L. grown in the drained paddy field and upland field. Grassland Science, 46: 133-142

Tobisa, M. ; Nakano, Y. ; Paek, J. ; Mochizuki, T. ; Shimojo, M. ; Masuda, Y., 2005. Fermentation quality and palatability of rice plant silages as affected by mixing with phasey bean (*Macroptilium lathyroides* (L.) Urb.). J. Japan. Soc. Grassl. Sci., 51 (3):274-280

Tobisa, M. ; Nakano, Y. ; Paek, J. ; Mochizuki, T. ; Shimojo, M. ; Masuda, Y., 2006. Fermentation quality of floating rice plant (*Oryza sativa* L.) silages as affected by mixing with phasey bean (*Macroptilium lathyroides* (L.) Urb.). J. Japan. Soc. Grassl. Sci., 52 (3):155-160

US Forest Service, 2012. *Macroptilium lathyroides* L. Pacific Island Ecosystems at Risk (PIER). Online resource

Werner, D. ; Newton, W. E., 2005. Nitrogen fixation in agriculture, forestry, ecology, and the environment: Volume 4. Springer

Whiteman, P. C. ; Seidlheko, M. ; Siregar, M. E., 1983. Comparative flooding tolerance of tropical pasture legumes. Proceedings of the XIV International Grassland Congress, held at Lexington, Kentucky, U.S.A., June 15-24, 1981 [Smith, J.A.; Hays, V.W. (Editors)]:393-395

Whiteman, P. C. ; Seidlheko, M. ; Siregar, M. E. ; Chudasama, A. K. ; Javier, R. R., 1984. Short-term flooding tolerance of seventeen commercial tropical pasture legumes. Trop. Grassl., 18 (2): 91-96

Yunus, M. ; Ohba, N. ; Tobisa, M. ; Nakano, Y. ; Shimojo, M. ; Furuse, M. ; Masuda, Y., 2001. Improving fermentation and nutritive quality of napiergrass silage by mixing with phasey bean. Asian-Aust. J. Anim. Sci., 14 (7): 947-950

Zhao, J. Y. ; Shimojo, M. ; Goto, I., 1993. The effects of feeding level and roughage/concentrate ratio on the measurement of protein degradability of two tropical forages in the rumen of goats, using the nylon bag technique. Anim. Feed Sci. Technol., 41 (4): 261-269

51 references found

Datasheet citation

Heuzé V., Tran G., Giger-Reverdin S., Bastianelli D., Lebas F., 2015. *Phasey bean (Macroptilium lathyroides)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/627>. Last updated on September 30, 2015, 17:30

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollot (AFZ)

Image credits

● Forest and Kim Starr ● Forest and Kim Starr

